

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 842 662 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
17.07.2002 Bulletin 2002/29

(51) Int Cl.⁷: **A61K 31/445**, A61K 9/70,
A61K 47/12

(21) Application number: **97918413.2**

(86) International application number:
PCT/JP97/01595

(22) Date of filing: **13.05.1997**

(87) International publication number:
WO 97/42952 (20.11.1997 Gazette 1997/50)

(54) PERCUTANEOUS TAPE PREPARATION CONTAINING FENTANYL

FENTANYL ENTHALTENDES TRANSDERMAL-PFLASTER

PREPARATION EN RUBAN PERCUTANEE CONTENANT DU FENTANYL

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**

- **SUZUKI, Tatsuaki, Tsukuba Lab. of Hisamitsu Tsukuba-shi, Ibaraki 305 (JP)**
- **HIGO, Naruhito, Tsukuba Lab. of Hisamitsu Tsukuba-shi, Ibaraki 305 (JP)**

(30) Priority: **13.05.1996 JP 14221096**

(43) Date of publication of application:
20.05.1998 Bulletin 1998/21

(74) Representative: **Reitzner, Bruno, Dr.
Splanemann Reitzner
Baronetzky Westendorp
Patentanwälte
Rumfordstrasse 7
80469 München (DE)**

(73) Proprietor: **HISAMITSU PHARMACEUTICAL CO.,
INC.
Tosu-shi Saga 841 (JP)**

(72) Inventors:
• **CHONO, Hideharu, Tsukuba Lab. of Hisamitsu Tsukuba-shi, Ibaraki 305 (JP)**
• **TERAHARA, Takaaki, Tsukuba Lab. of Hisamitsu Tsukuba-shi, Ibaraki 305 (JP)**

(56) References cited:
EP-A- 0 757 910 EP-A- 0 788 792
WO-A-89/10108 JP-A- 4 099 719
JP-A- 4 321 624 JP-A- 6 040 947
JP-A- 62 126 119 US-A- 5 186 939

EP 0 842 662 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention relates to a tape formulation for percutaneous administration containing fentanyl (chemical name: 1-phenethyl-4-N-propionyl-anilino-piperidine) or a salt thereof, which is very excellent in transdermal permeable property and which has a low irritative property to skins. The tape formulation for percutaneous administration containing fentanyl of the present invention is greatly expected to be utilized as a prolonged-action anesthetic and analgesic.

Background Art

[0002] Fentanyl, in particular fentanyl citrate is known as a pharmaceutical having a high analgesic effect. However, there was no useful administration method of the pharmaceutical for relatively long lasting pains such as carcinomatous pains, since its elimination half life is short and thus its effects does not last though it is utilized for the constant rate instillation before and after operation.

[0003] In the USA, a prolonged-action patch formulation containing fentanyl base (trade name: DURAGESIC) is put on the market. However, it has the disadvantage of being highly irritative to administered regions (The PII value showing a primary irritant index to rabbit skins of the patch formulation is 2.2, which is a very high value compared with that of the formulation of the present invention which is 0.3 to 0.8 (see Table 3).

[0004] Further, although the attempts to formulate fentanyl citrate into a tape formulation for percutaneous administration have been made, it could not be used clinically, since the solubility of fentanyl citrate in nonaqueous base is low and thus the transdermal permeable property of the formulation, in which fentanyl is contained in a nonaqueous base, is very low.

[0005] Therefore, an object of the present invention is to dissolve the above problems in connection with the prior art, and to provide a tape formulation for percutaneous administration containing fentanyl, which has a low irritative property to skin, which is extremely excellent in the transdermal permeable property of fentanyl, and which is stable during the storage period.

[0006] The present inventors have researched earnestly in order to achieve the above object, and as a result, they found that a tape formulation for percutaneous administration which is extremely excellent in transdermal permeable property and which has a low irritative property to skin, can be prepared, by adding sodium acetate to a pressure sensitive adhesive base containing fentanyl or a salt thereof, and thus completed the present invention.

[0007] EP-A-788 792 (prior art in the sense of Art. 54(3) (4)) discloses matrix type patch (i.e. tape) formulations for percutaneous administration of a physiologically active substance, e.g. fentanyl (citrate). Examples 2 and 16 describe formulations comprising a pressure sensitive adhesive, fentanyl citrate and sodium acetate. The formulations are as follows:

	Example 2	Example 16
- SIS* (Califlex® D-1111; Shell Co)	16,5 %	-
- PIB** (Vistanex; MH-L-100; Exxon Co.)	1,5 %	15 %
- PIB** (Vistanex LMMH); Exxon Co.)	6,0 %	-
- Alicyclic saturated hydrocarbon resin (Arcon® P-199; Arkawa Kagaku Co. Ltd.)	30 %	-
- Liquid paraffin (Crystal® 352, Esso Co.)	37 %	-
- Pirodiodecane	3 %	2,5 %
- Sodium acetate	1,5 %	0,5 %
- Fentanyl citrate	3 %	2 %
- Aluminum silicate	1 %	-
- BHT*** (Yoshinox®; Yoshitomi Pharm. Ind. Co.)	0,5%	-
- Hardened rosin ester (Foral® 85; Rike Hercules Co.)	-	70 %
- Crotomiton	-	10 %
Total Amount	100 %	100 %

*SIS = styrene-isoprene-styrene block copolymer

**PIB = polyisobutylene

***BHT = dibutyl hydroxytoluene

Disclosure of Invention

[0008] Thus the present invention relates to a tape formulation for percutaneous administration containing fentanyl which comprises fentanyl or a salt thereof, a pressure sensitive adhesive and sodium acetate, except the formulations of Examples 2 and 16 of EP-A-788 792 referred to above.

[0009] The present invention also relates to the tape formulation for percutaneous administration containing fentanyl, which comprises 0.05 to 20 % (w/w) of fentanyl or a salt thereof, 0.1 to 98 % (w/w) of a pressure sensitive adhesive and 0.01 to 15 % (w/w) of sodium acetate.

[0010] The present invention also relates to the tape formulation for percutaneous administration containing fentanyl, wherein the salt of fentanyl is fentanyl citrate.

[0011] The present invention also relates to the tape formulation for percutaneous administration containing fentanyl, wherein the weight ratio of formulation of fentanyl citrate and sodium acetate is (1 to 5) : (0.5 to 2.5).

[0012] The present invention also relates to the tape formulation for percutaneous administration containing fentanyl, wherein the weight ratio of formulation of fentanyl citrate and sodium acetate is (3 to 5) : (1.5 to 2.5).

[0013] The present invention also relates to the tape formulation for percutaneous administration containing fentanyl, wherein the weight ratio of formulation of fentanyl citrate and sodium acetate is 2 : 1.

[0014] The present invention also relates to the tape formulation for percutaneous administration containing fentanyl, which further comprises an oil and/or a tackifier.

[0015] The present invention also relates to the tape formulation for percutaneous administration containing fentanyl, which further comprises a transdermal absorption enhancer.

[0016] The present invention also relates to the tape formulation for percutaneous administration containing fentanyl, wherein the pressure sensitive adhesive comprises two components of polyisobutylene and styrene-isoprene-styrene block copolymer.

Best Mode For Carrying Out The Invention

[0017] The tape formulation for percutaneous administration containing fentanyl of the present invention will be explained in detail, hereinafter.

[0018] The pharmacological active component of the tape formulation for percutaneous administration containing fentanyl of the present invention, is fentanyl itself or a salt thereof. The salt of fentanyl is not particularly limited, and an inorganic salt and an organic salt thereof may be used. As typical fentanyl salts, citrate, hydrochloride, fumarate and the like may be exemplified. Among them, fentanyl citrate is particularly preferable. The fentanyl or a salt thereof may be used alone, and a mixture of at least two of them may be used.

[0019] The fentanyl or a salt thereof is preferably contained in an amount ranging from 0.05 to 20 % (w/w) based on the total amount of the adhesive layer of the tape formulation for percutaneous administration of the present invention. If the amount of fentanyl or a salt thereof is less than 0.05 % (w/w), a sufficient permeation amount can not be obtained as a tape formulation for percutaneous administration, and if the amount exceeds 20 % (w/w), it exerts a bad influence on the physical properties of the formulation itself and thus it is not preferable.

[0020] The pressure sensitive adhesive contained in the adhesive layer of the tape formulation for percutaneous administration containing fentanyl of the present invention, is not limited, but polyisobutylene (PIB), styrene-isoprene-styrene block copolymer (SIS) [e.g., Califlex D-1111, Califlex Tr-1107 manufactured by Shell Chemical; JSR5000, JSR-5002, SR5100 manufactured by Japan Synthetic Rubber Co., Ltd.; Quintack 3421 manufactured by Nippon Zeon Co., Ltd.], isoprene rubber, styrene-butadiene-styrene copolymer (SBS) [e.g., Califlex TR-1101 manufactured by Shell Chemical], acrylic polymer [e.g., a copolymer comprising at least two components selected from the group comprising 2-ethylhexyl acrylate, vinyl acetate, ethyl acrylate, methacrylate, methoxyethyl acrylate and acrylic acid, such as PE-300 (manufactured by Nippon Carbide Industries Co., Inc.)] may be exemplified as preferable examples. These polymers may be used alone or a mixture of at least two of them may be used. Among them, two components comprising PIB and SIS is preferably used. In this case, the weight ratio of formulation of PIB and SIS is preferably 1:1 to 1:4.

[0021] The pressure sensitive adhesive is preferably contained in an amount ranging from 0.1 to 98 % (w/w), more preferably from 0.1 to 70 % (w/w), most preferably from 0.1 to 50 % (w/w), based on the total amount of the adhesive layer of the tape formulation for percutaneous administration of the present invention. If the amount of the pressure sensitive adhesive is less than 0.1 % (w/w), the physical properties of the formulation itself will be poor and thus it is not preferable. If the amount exceeds 98 % (w/w), a satisfactory adhesive property to human skin can not be obtained and it is not preferable.

[0022] The transdermal permeable property of fentanyl or a salt thereof is highly increased by formulating sodium acetate in the adhesive layer of the tape formulation for percutaneous administration containing fentanyl of the present invention. The sodium acetate is preferably contained in an amount ranging from 0.01 to 15 % (w/w), more preferably from 0.01 to 10 % (w/w), most preferably from 0.01 to 5 % (w/w), based on the total amount of the adhesive layer. If the

amount of sodium acetate is less than 0.01 % (w/w), the effect of remarkably improving the transdermal permeable property can not be obtained. If the amount exceeds 15 % (w/w), the irritativeness to skin is increased, thus it is not preferable.

[0023] If the fentanyl salt is fentanyl citrate, the maximum effects may be obtained in the respects of the physical property and transdermal permeable property when the weight ratio of formulation of fentanyl citrate and sodium acetate is (1 to 5) : (0.5 to 2.5), preferably (3 to 5) : (1.5 to 2.5), more preferably 2 : 1. The amount of sodium acetate is less than the ratio of formulation, the transdermal permeable property of the drug is decreased suddenly, and the amount of sodium acetate exceeds the ratio of formulation, the tape formulation will be inhomogeneous and the physical properties such as adhesive property will be poor and thus it is not preferable.

[0024] In addition, a tackifier may be formulated in the adhesive layer of the formulation of the present invention so as to impart an adhesive property to the formulation since the adhesive property of the pressure sensitive adhesive is low. As preferable examples of the tackifiers, polyterpene resins, petroleum resins, rosins, rosin esters, oil-soluble phenol resins and the like may be exemplified. As the concrete examples of the tackifiers, Clearon P-105, Foral 105, Arcon P-100, KE-311, KE-100, Super Ester S-100, Tamanol 521, YS Resin 75, KR-610 and the like may be exemplified by trade names.

[0025] The tackifiers is preferably contained in the range from 0.1 to 70 % (w/w), more preferably from 5 to 50 % (w/w), and most preferably from 10 to 35 % (w/w), based on the total amount of the adhesive layer of the formulation of the present invention.

[0026] Further, an oil may be formulated in the adhesive layer as a softening agent in order to improve the processability and to control the adhesive property of the tape formulation for percutaneous administration of the present invention. As the oil, for example, liquid paraffin, squalane, olive oil, Tsubaki oil, Persic oil and peanut oil are preferable, and liquid paraffin is most preferable.

[0027] The oil is preferably contained in an amount ranging from 1 to 70 % (w/w), more preferably from 10 to 60 % (w/w), and most preferably from 20 to 50 % (w/w), based on the total amount of the adhesive layer of the formulation of the present invention.

[0028] Further, a transdermal absorption enhancer may be formulated in the adhesive layer of the formulation of the present invention depending on the necessities. As the transdermal absorption enhancer, any compounds an absorption enhancing effect in skins of which is recognized, may be used. For example, fatty acids having carbon chains of 6 to 20, aliphatic alcohols, fatty acid esters or ethers, aromatic organic acids, aromatic alcohols, aromatic organic acid esters or ethers may be exemplified. In addition, lactic acid esters, acetic acid esters, monoterpene compounds, sesquiterpene compounds, Azone or its derivatives, glycerin fatty acid esters, sorbitan fatty esters, polysorbates, polyethylene glycol fatty acid esters, polyoxyethylene hardened castor oils, sucrose fatty acid esters and the like may be exemplified. As the concrete examples of the enhancer, caprylic acid, capric acid, caproic acid, lauric acid, myristic acid, palmitic acid, stearic acid, oleic acid, linolic acid, linolenic acid, lauryl alcohol, myristyl alcohol, oleyl alcohol, cetyl alcohol, methyl laurate, isopropyl myristate, myristyl myristate, octyldodecyl myristate, cetyl palmitate, salicylic acid, methyl salicylate, ethylene glycol salicylate, cinnamic acid, methyl cinnamate, cresol, cetyl lactate, ethyl acetate, propyl acetate, geraniol, thymol, eugenol, terpineol, l-menthol, borneol, d-limonene, isoeugenol, isoborneol, nerol, dl-camphor, glycerol monolaurate, glycerol monooleate, sorbitan monolaurate, sucrose monolaurate, polysorbate 20, polyethylene glycol monolaurate, polyethylene glycol monostearate, HCO-60 (hardened castor oil), 1-[2-(decyl thio)ethyl] azacyclopentane-2-one (hereinafter it abbreviated as "pirotiodecane") are preferable, and lauryl alcohol, myristyl alcohol, ethylene glycol salicylate and pirotiodecane are most preferable.

[0029] These transdermal absorption enhancers is preferably contained in an amount ranging from 0.01 to 20 % (w/w), more preferably in an amount ranging from 0.1 to 10 % (w/w), most preferably in an amount ranging from 0.5 to 5 % (w/w), based on the total amount of the adhesive layer of the tape of the present invention. If the amount of the transdermal absorption enhancer exceeds 20 % (w/w), the irritations to skin such as erythema and edema are shown, and if the amount is less than 0.01 % (w/w), the effect of formulating the transdermal absorption enhancer can not be obtained, and thus it is not preferable.

[0030] In addition, a hydrophilic polymer may be contained in the tape formulation of the present invention depending on the necessities in order to absorb the aqueous components such as sweat produced from skins. As the hydrophilic polymer, for example, light silicic anhydride, cellulose derivatives [carboxymethylcellulose (CMC), sodium carboxymethylcellulose (CMCNa), methylcellulose (MC), hydroxypropyl methylcellulose (HPMC), hydroxypropyl cellulose (HPC), hydroxyethyl cellulose (HEC)], starch derivatives (Pullulan), polyvinyl alcohol (PVA), polyvinyl pyrrolidone (PVP), vinyl acetate (VA), carboxyvinyl polymer (CVP), ethyl vinyl acetate (EVA), Eudragit, gelatin, polyacrylic acid, sodium polyacrylate, polyisobutylene maleic anhydride copolymer, alginic acid, sodium alginate, carrageenan, gum arabi, tragacanth, gum karaya, and polyvinyl methacrylate are preferable, and light silicic anhydride, cellulose derivatives (CMCNa, HPMC, HPC, MC) and Eudragit are more preferable.

[0031] The hydrophilic polymer is preferably contained in an amount ranging from 0.1 to 20 % (w/w), more preferably 0.5 to 10 % (w/w), based on the total amount of the adhesive layer of the tape formulation for percutaneous adminis-

tration of the present invention.

[0032] In addition, a cross-linking agent, a preservative, an antioxidant and another components may be formulated in the adhesive layer of the formulation of the present invention.

[0033] As the cross-linking agent, a thermosetting resin such as an amino resin, a phenol resin, an epoxy resin, an alkyd resin, an unsaturated polyester, an isocyanate compound, a block isocyanate compound, an organic cross-linking agent, and an inorganic corss-linking agent such as a metal or a metal compound are preferable. As the preservative, ethyl p-oxy benzoate, propyl p-oxy benzoate, butyl p-oxy benzoate and the like are preferable. As the antioxidant, tocopherol and its ester derivatives, ascorbic acid, stearic acid ester, nordihydroguaiaretic acid, dibutyl hydroxytoluene (BHT), butyl hydroxy anisol (BHA) and the like are preferable.

[0034] The adhesive layer of the tape formulation of the present invention preferably comprises a nonaqueous base, and the effects of the present invention may be obtained effectively by making the base nonaqueous.

[0035] The adhesive layer comprising the above comonents may be prepared by any conventional methods. For example, when the layer is prepared by a solvent method, the formulation of the present invention may be prepared by adding other components other than polymers to an organic solvent solution of the polymers, then stirring, and applying the mixture on a backing film and drying. When the polymers to be formulated can be applied by a hot-melt method, the formulation of the present invention may be obtained by dissolving the polymer components at a high temperature, then adding another components, stirring, and applying on a backing film.

[0036] The tape formulation of the present invention may comprise any layers provided that it has an adhesive layer having the above components, and another layers and components of the layers are not limited. For example, the tape formulation for percutaneous administration may comprise, besides the adhesive layer, a backing layer supporting the adhesive layer, a releasable liner layer provided on the adhesive layer, and the like.

[0037] The backing layer may comprise, for example, fabric, non-woven fabric, polyurethane, polyvinyl acetate, polyvinylidene chloride, polyethylene, polyethylene terephthalate, polybutylene terephthalate, paper, aluminum sheet and the like, and a composite material thereof.

[0038] The tape formulation for percutaneous administration of the present invention is a very useful means for mitigating pains for the patients who can not take orally a anesthetic analgesic easily, since fentanyl or a salt thereof is absorbed continuously via skins by the tape formulation of the present invention. In addition, the tape formulation of the present invention can be administered non-invasively and thus can decrease the burdens of the patients compared with the continuous intradermal aministration method being an invasive method. Further, as to the dose can be controlled easily, e.g., by cutting the formulation, depending on the conditions, ages, body weights, sex distinctions and another factors of patients.

Examples

[0039] The present invention will be explained in more detail with the following Examples, hereinafter. However, the present invention is not be limited to the Examples, and the present invention extends to all such modifications and variations as will be apparent to those skilled in the art without departing the scope of the present invention. In the Examples, all "%" are %(w/w) unless noted.

Example 1

[0040]

Sodium acetate	2.5%
Acrylic polymer(PE-300)	88.5%
Toluene diisocyanate	1.0%
Pirotiodecane	3.0%
Fentanyl citrate	5.0%
-----	-----
Total Amount	100%

[0041] Sodium acetate, pirotiodecane and fentanyl citrate were added to ethanol, and stirred to dissolve them at room temperature. Then, a solution of acrylic polymer in ethyl acetate and toluene diisocyanate were added to the mixture and stirred. The mixture was applied onto a polyethylene terephthalate film (PET) (30µm), and it was cross-linked thermally at 90°C for 15 minutes so as to have an adhesive layer of 50µm. Using the adhesive layer, a tape formulation for percutaneous administration of the present invention was prepared by a conventional method.

Example 2**[0042]**

5	Sodium acetate	1.5%
	Pirotiodecane	3.0%
	Liquid paraffin	38.0%
	Polyterpene resin tackifier	29.5%
10	Polyisobutylene	7.5%
	Styrene-isoprene-Styrene block copolymer	16.5%
	Antioxidant (BHT)	0.5%
	Aluminum silicate	0.5%
15	Fentanyl citrate	3.0%

	Total Amount	100%

20 **[0043]** After the components except sodium acetate, pirotiodecane and fentanyl citrate were dissolved and mixed at 180°C, the rest components were added and dispersed so as to have a homogeneous mixture. Then the mixture was applied onto a PET film (30µm) to have an adhesive layer of 100µm. Using the adhesive layer, a tape formulation for percutaneous administration of the present invention was prepared by a conventional method.

Example 3**[0044]**

25	Sodium acetate	2.5%
	Pirotiodecane	3.0%
	Liquid paraffin	39.5%
30	Polyterpene resin tackifier	21.7%
	Polyisobutylene	6.8%
	Styrene-isoprene-styrene block copolymer	20.4%
	Antioxidant (BHT)	0.5%
35	Aluminum silicate	0.6%
	Fentanyl citrate	5.0%

	Total Amount	100%

40 **[0045]** After the components except sodium acetate, pirotiodecane and fentanyl citrate were dissolved and mixed at 180°C, the rest components were added and dispersed so as to have a homogeneous mixture. Then the mixture was applied onto a PET film (30µm) to have an adhesive layer of 100µm. Using the adhesive layer, a tape formulation for percutaneous administration of the present invention was prepared by a conventional method.

Example 4**[0046]**

50	Sodium acetate	2.5%
	Liquid paraffin	12.5%
	Oil-soluble phenol resin tackifier	39.5%
	Polyisobutylene	7.5%
	Styrene-isoprene-styrene block copolymer	30.5%
	Antioxidant (BHT)	0.5%
55	Lauryl alcohol	2.0%
	Fentanyl citrate	5.0%

	Total Amount	100%

[0047] After the components except lauryl alcohol, sodium acetate and fentanyl citrate were dissolved and mixed at 180°C, the rest components were added and dispersed so as to have a homogeneous mixture. Then the mixture was applied onto a PET film (30µm) to have an adhesive layer of 100µm. Then using the adhesive layer, a tape formulation for percutaneous administration of the present invention was prepared by a conventional method.

5
Example 5

[0048]

Sodium acetate	1.5%
Crotamiton	3.0%
Liquid paraffin	38.5%
Polyterpene resin tackifier	29.5%
Polyisobutylene	7.5%
Styrene-isoprene-styrene block copolymer	16.5%
Antioxidant (BHT)	0.5%
Fentanyl citrate	3.0%

Total Amount	100%

[0049] After stirring to dissolve sodium acetate, crotamiton, fentanyl citrate and liquid paraffin at 80°C, the mixture was mixed with a cyclohexane solution in which styrene-isoprene-styrene block copolymer, polyisobutylene, polyterpene resin tackifier and antioxidant had been dissolved previously. Then the mixture was applied onto a PET film (30µm), and dried at 85°C for 30 minutes so as to have an adhesive layer of 50µm. Using the adhesive layer, a tape formulation for percutaneous administration of the present invention was prepared by a conventional method.

Example 6

[0050]

Sodium acetate	2.5%
Liquid paraffin	35.0%
Polyterpene resin tackifier	25.5%
Polyisobutylene	7.0%
Styrene-isoprene-styrene block copolymer	24.0%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl	5.0%

Total Amount	100%

[0051] After the components except sodium acetate and fentanyl were dissolved and mixed at 180°C, the rest components were added and dispersed so as to have a homogeneous mixture. Then the mixture was applied onto a PET film (30µm) to have an adhesive layer of 100µm. Using the adhesive layer, a tape formulation for percutaneous administration of the present invention was prepared by a conventional method.

Example 7

[0052]

Sodium acetate	0.5%
Pirotidecane	3.0%
Liquid paraffin	29.0%
Polyterpene resin tackifier	42.1%
Polyisobutylene	7.0%
Styrene-isoprene-styrene block copolymer	16.4%

EP 0 842 662 B1

(continued)

Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	1.0%

Total Amount	100%

[0053] Using the above components (Fentanyl citrate : Sodium acetate = 2:1), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

Example 8

[0054]

Sodium acetate	1.5%
Pirotidecane	3.0%
Liquid paraffin	28.8%
Polyterpene resin tackifier	41.5%
Polyisobutylene	6.9%
Styrene-isoprene-styrene block copolymer	16.2%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	1.0%

Total Amount	100%

[0055] Using the above components (Fentanyl citrate : Sodium acetate = 2:3), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

Example 9

[0056]

Sodium acetate	2.5%
Pirotidecane	3.0%
Liquid paraffin	28.7%
Polyterpene resin tackifier	41.0%
Polyisobutylene	6.8%
Styrene-isoprene-styrene block copolymer	16.0%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	1.0%

Total Amount	100%

[0057] Using the above components (Fentanyl citrate : Sodium acetate = 2:5), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

Example 10

[0058]

Sodium acetate	0.5%
Pirotidecane	3.0%
Liquid paraffin	28.7%

(continued)

Polyterpene resin tackifier	41.0%
Polyisobutylene	6.8%
Styrene-isoprene-styrene block copolymer	16.0%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	3.0%

Total Amount	100%

[0059] Using the above components (Fentanyl citrate : Sodium acetate = 6:1), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

Example 11

[0060]

Sodium acetate	1.5%
Pirotidecane	3.0%
Liquid paraffin	28.5%
Polyterpene resin tackifier	40.5%
Polyisobutylene	6.8%
Styrene-isoprene-styrene block copolymer	15.7%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	3.0%

Total Amount	100%

[0061] Using the above components (Fentanyl citrate : Sodium acetate = 2:1), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

Example 12

[0062]

Sodium acetate	2.5%
Pirotidecane	3.0%
Liquid paraffin	28.2%
Polyterpene resin tackifier	40.0%
Polyisobutylene	6.7%
Styrene-isoprene-styrene block copolymer	15.6%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	3.0%

Total Amount	100%

[0063] Using the above components (Fentanyl citrate : Sodium acetate = 6:5), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

Example 13

[0064]

5

Sodium acetate	0.5%
Pirotiodecane	3.0%
Liquid paraffin	28.2%
Polyterpene resin tackifier	40.0%
Polyisobutylene	6.7%
Styrene-isoprene-styrene block copolymer	15.6%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	5.0%

Total Amount	100%

10

15

[0065] Using the above components (Fentanyl citrate : Sodium acetate = 10:1), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

20

Example 14

[0066]

25

Sodium acetate	1.5%
Pirotiodecane	3.0%
Liquid paraffin	28.2%
Polyterpene resin tackifier	39.5%
Polyisobutylene	6.5%
Styrene-isoprene-styrene block copolymer	15.3%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	5.0%

Total Amount	100%

30

35

[0067] Using the above components (Fentanyl citrate : Sodium acetate = 10:3), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

40

Example 15

[0068]

45

Sodium acetate	2.5%
Pirotiodecane	3.0%
Liquid paraffin	28.0%
Polyterpene resin tackifier	38.9%
Polyisobutylene	6.5%
Styrene-isoprene-styrene block copolymer	15.1%
Antioxidant (BHT)	0.5%
Aluminum silicate	0.5%
Fentanyl citrate	5.0%

Total Amount	100%

50

55

[0069] Using the above components (Fentanyl citrate : Sodium acetate = 2:1), a tape formulation for percutaneous administration of the present invention was prepared by the method as described in Example 2.

Comparative Examples 1 to 5

[0070] Comparative Examples 1 to 5 are each corresponding to Examples 1 to 5, respectively. In each Comparative Example, a tape formulation for percutaneous administration was prepared by the method as described in the corresponding Example, provided that sodium acetate which was used in Examples 1 to 5, was not formulated.

Test Example 1

(In vitro transdermal permeation test)

[0071] As to each of the formulations for percutaneous administration obtained in Examples 1 to 5, 7 to 15 and Comparative Examples 1 to 5, the evaluations were made by an in vitro transdermal permeation test using hairless mouse skins.

[0072] After picking out the skin of the back region of hairless mouse (6 to 9 weeks old), fats in the dermal side were removed carefully. The skin was installed in a flow-through cell, in which water at 37° C was circulated around the outer periphery of the receptor layer, so that the dermal side is to be a receptor layer. Each of the tape formulations for percutaneous administration obtained in Examples 1 to 5, 7 to 15 and Comparative Examples 1 to 5 was applied onto the stratum corneum side, and samplings were made at every one hour for 24 hours at a rate of 5ml/hour using physiological saline to the receptor layer. Then, the flow rates at every one hour were determined accurately, and the drug concentrations were determined by a high-performance liquid chromatography method. The permeation rates at one hour were calculated according to the following formula, and the permeation rates in steady state were determined. The results are shown in Table 1.

$$\text{Transdermal Permeation Rate } (\mu\text{g}/\text{cm}^2/\text{hr}) = [\text{Drug concentration } (\mu\text{g}/\text{ml}) \times \text{Flow Rate (ml)}] / \text{Applied surface area of the formulation (cm}^2\text{)}$$

Table 1

	Transdermal Permeation Rate ($\mu\text{g}/\text{cm}^2/\text{hr}$)
Example 1	15.5
Example 2	25.3
Example 3	36.8
Example 4	35.2
Example 5	22.3
Example 7	8.8
Example 8	8.2
Example 9	7.6
Example 10	9.4
Example 11	22.2
Example 12	20.5
Example 13	12.2
Example 14	29.4
Example 15	35.8
Comparative Example 1	1.2
Comparative Example 2	1.0
Comparative Example 3	1.2
Comparative Example 4	1.5
Comparative Example 5	1.1

[0073] As it is clear from Table 2, the tape formulations for percutaneous administration obtained in Examples 1 to

5 and 7 to 15 have higher transdermal permeation rates, compared with the tape formulations for percutaneous administration obtained in Comparative Examples 1 to 5.

[0074] In particular, it is proved that the tape formulations for percutaneous administration of Examples 1 to 5, 7, 11, 12, 14 and 15, in which the formulation ratios of fentanyl citrate and sodium acetate are (3 to 5) : (1.5 to 2.5), have very high transdermal permeation rates.

[0075] Among them, it is proved that the tape formulations for percutaneous administration of Examples 1 to 5, 11 and 15, in which the formulation ratios of fentanyl citrate and sodium acetate are 2 : 1, have extremely high transdermal permeation rates.

Test Example 2

(Primary irritant to rabbit skin test)

[0076] As to each of the tape formulations for percutaneous administration obtained in Examples 1 to 5, the evaluations were made by an in vivo primary irritant test using a rabbit skin.

[0077] Each of the tape formulations for percutaneous administration obtained in Examples 1 to 5 was applied onto rabbit skin. The decisions were made in accordance with the criterion of irritativeness to skin shown in Table 2, as to the erythema and edema at 24 and 48 hours after the application. The total of the both scores were regarded as the irritative score at each time. Further, the average value of the irritative scores at each time was regarded as a PII Value. In addition, as the control groups, patch tape of Pharmacopoeia Japonica and a commercially available product in USA (DURAGESIC) were used. The results are shown in Table 3.

Table 2

Criterion of Irritiveness to Skin		
Score	Erythema	Edema
0	None	None
1	Extremely slight	Extremely slight
2	Evident	Evident
3	Middle degree up to intense	Middle degree up to intense
4	Scarlet, intense	Intense

Table 3

	Irritaviness to Skin (PII Value)
Example 1	0.5
Example 2	0.7
Example 3	0.5
Example 4	0.7
Example 5	0.3
Adhesive tape of Pharmacopoeia Japonica	0.3
DURAGESIC (commercially available in USA)	2.2*

*: extracts from the application documents to FDA

[0078] From the results shown in Table 2, it is proved that the tape formulations for percutaneous administration of Examples 1 to 5 have very low irritative property to skins compared with the conventional product (DURAGESIC), and have the equal irritative property to skins to the patch tape of Pharmacopoeia Japonica which has low irritative property.

Industrial Applicability

[0079] With the present invention, fentanyl of a salt thereof can be formulated into a formulation for percutaneous administration which is low-irritative and excellent in transdermal permeable property, which could not be attained by the prior arts.

[0080] Thus fentanyl or a salt thereof can be delivered into the body and the pharmacological effects of fentanyl or

a salt thereof can be utilized effectively and continuously, by using the tape formulation for percutaneous administration containing fentanyl of the present invention.

[0081] Therefore, the tape formulation for percutaneous administration containing fentanyl of the present invention will be a very useful means for mitigating pains for the patients who can not take orally an anesthetic analgesic easily.

5

Claims

1. A tape formulation for percutaneous administration containing fentanyl, which comprises fentanyl or a salt thereof, a pressure sensitive adhesive and sodium acetate, except the following formulations:

10

- SIS* (Califlex® D-1111; Shell Co)	16,5 %	-
- PIB**(Vistanex; MH-L-100; Exxon Co.)	1,5 %	15 %
- PIB**(Vistanex LMMH); Exxon Co.)	6,0 %	-
- Alicyclic saturated hydrocarbon resin (Arcon® P-199; Arkawa Kagaku Co. Ltd.)	30 %	-
- Liquid paraffin (Crystal® 352, Esso Co.)	37 %	-
- Pirotiodecane	3 %	2,5%
- Sodium acetate	1,5 %	0,5 %
- Fentanyl citrate	3 %	2 %
- Aluminum silicate	1%	-
- BHT*** (Yoshinox®; Yoshitomi Pharm. Ind. Co.)	0,5%	-
- Hardened rosin ester (Foral® 85; Rike Hercules Co.)	-	70 %
- Crotomiton	-	10 %
Total Amount	100 %	100 %

15

20

25

*SIS = styrene-isoprene-styrene block copolymer

**PIB = polyisobutylene

***BHT = dibutyl hydroxytoluene

30

2. A tape formulation for percutaneous administration containing fentanyl of Claim 1, which comprises 0.05 to 20 % (w/w) of fentanyl or a salt thereof, 0.1 to 98 % (w/w) of a pressure sensitive adhesive and 0.01 to 15 % (w/w) of sodium acetate.
3. A tape formulation for percutaneous administration containing fentanyl of Claim 1 or 2, wherein the fentanyl salt is fentanyl citrate.
4. A tape formulation for percutaneous administration containing fentanyl of Claim 3, wherein the weight ratio of formulation of fentanyl citrate and sodium acetate is (1 to 5) : (0.5 to 2.5).
5. A tape formulation for percutaneous administration containing fentanyl of Claim 3, wherein the weight ratio of formulation of fentanyl citrate and sodium acetate is (3 to 5) : (1.5 to 2.5).
6. A tape formulation for percutaneous administration containing fentanyl of Claim 3, wherein the weight ratio of formulation of fentanyl citrate and sodium acetate is 2 : 1.
7. A tape formulation for percutaneous administration containing fentanyl of any of Claims 1 to 6, which further comprises an oil and/or a tackifier.
8. A tape formulation for percutaneous administration containing fentanyl of any of Claims 1 to 7, which further comprises a transdermal absorption enhancer.
9. A tape formulation for percutaneous administration containing fentanyl of any of Claims 1 to 8, wherein the pressure sensitive adhesive comprises two components of polyisobutylene and styrene-isoprene-styrene block copolymer.

55

Patentansprüche

1. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, welche Fentanyl oder ein Salz davon, einen druckempfindlichen Klebstoff und Natriumacetat enthält, ausgenommen die folgenden Formulierungen:

5

10

15

20

- SIS* (Califlex® D-1111; Shell Co.)	16,5%	-
- PIB** (Vistanex; MH-L-100; Exxon Co.)	1,5%	15%
- PIB** Vistanex LMMH; Exxon Co.)	6,0%	-
- Alicyclisches gesättigtes Kohlenwasserstoffharz (Arcon® P-199; Arkawa Kagaaku Co. Ltd.)	30%	-
- Flüssiges Paraffin (Cristol® 352, Esso Co.)	37%	-
- Pirotiodecan	3%	2,5%
- Natriumacetat	1,5%	0,5%
- Fentanylcitrat	3%	2%
- Aluminumsilicat	1%	1%
- BHT*** (Yoshinox®; Yoshitomi Pharm. Ind. Co.)	0,5%	-
- Gehärteter Rosinester (Foral® 85; Rike Hercules Co.)	-	70%
- Crotomiton	-	10%
Gesamtbetrag	100%	100%

*SIS = Styrol-Isopren-Styrol Block-Copolymer

**PIB = Polyisobutylene

***BHT = Dibutylhydroxytoluol

25

2. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, nach Anspruch 1, welche 0,05 bis 20% (w/w) Fentanyl oder ein Salz davon, 0,1 bis 98% (w/w) eines druckempfindlichen Klebstoffs und 0,01 bis 15% (w/w) Natriumacetat enthält.

30

3. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, nach Anspruch 1 oder 2, worin das Fentanylsalz Fentanylcitrat darstellt.

4. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, nach Anspruch 3, worin das Gewichtsverhältnis der Formulierung zwischen Fentanylcitrat und Natriumacetat (1 bis 5) : (0,5 bis 2,5) beträgt.

35

5. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, nach Anspruch 3, worin das Gewichtsverhältnis der Formulierung zwischen Fentanylcitrat und Natriumacetat (3 bis 5) : (1,5 bis 2,5) beträgt.

6. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, nach Anspruch 3, worin das Gewichtsverhältnis der Formulierung zwischen Fentanylcitrat und Natriumacetat 2 : 1 beträgt.

40

7. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, nach einem der Ansprüche 1 bis 6, welche zusätzlich ein Öl und/oder einen Klebrigmacher enthält.

8. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, nach einem der Ansprüche 1 bis 7, welche zusätzlich ein Mittel zur Förderung der transdermalen Absorption enthält.

45

9. Bandformulierung für die perkutane Anwendung, enthaltend Fentanyl, nach einem der Ansprüche 1 bis 8, worin der druckempfindliche Klebstoff zwei Komponenten, nämlich Polyisobutylene und Styrol-Isopren-Styrol-Blockcopolymer enthält.

50

Revendications

1. Formulation de bande pour administration percutanée contenant du fentanyl, qui comprend du fentanyl ou un sel de celui-ci, un adhésif autocollant et de l'acétate de sodium, à l'exception des formulations suivantes :

55

EP 0 842 662 B1

	SIS* (Califlex® D-1111 ; Shell Co.)	16,5 %	-
	PIB** (Vistanex ; MH-L-100 ; Exxon Co.)	1,5 %	15 %
5	PIB** (Vistanex LMMH ; Exxon Co.)	6,0 %	-
	Résine hydrocarbonée alicyclique saturée (Arcon® P-199 ; Arkawa Kagaku Co. Ltd.)	30 %	-
	Paraffine liquide (Cristol® 352, Esso Co.)	37 %	-
	Pirotiodécane	3 %	2,5 %
10	Acétate de sodium	1,5 %	0,5 %
	Citrate de fentanyl	3 %	2 %
	Silicate d'aluminium	1 %	-
15	BHT*** (Yoshinox® ; Yoshitomi Pharm. Ind. Co.)	0,5 %	-
	Ester de rosine durci (Foral® 85 ; Rike Hercules Co.)	-	70 %
	Crotomiton	-	10 %
	Quantité totale	100 %	100 %

* SIS = copolymère bloc styrène-isoprène-styrène

** PIB = polyisobutylène

***BHT = dibutylhydroxytoluène

2. Formulation de bande pour administration percutanée contenant du fentanyl selon la revendication 1, qui comprend 0,05 à 20 % (m/m) de fentanyl ou un sel de celui-ci, 0,1 à 98 % (m/m) d'un adhésif autocollant et 0,01 à 15 % (m/m) d'acétate de sodium.
3. Formulation de bande pour administration percutanée contenant du fentanyl selon la revendication 1 ou 2, dans laquelle le sel de fentanyl est le citrate de fentanyl.
4. Formulation de bande pour administration percutanée contenant du fentanyl selon la revendication 3, dans laquelle le rapport en poids dans la formulation de citrate de fentanyl et d'acétate de sodium est (1 à 5) : (0,5 à 2,5).
5. Formulation de bande pour administration percutanée contenant du fentanyl selon la revendication 3, dans laquelle le rapport en poids dans la formulation de citrate de fentanyl et d'acétate de sodium est (3 à 5) : (1,5 à 2,5).
6. Formulation de bande pour administration percutanée contenant du fentanyl selon la revendication 3, dans laquelle le rapport en poids dans la formulation de citrate de fentanyl et d'acétate de sodium est 2 : 1.
7. Formulation de bande pour administration percutanée contenant du fentanyl selon l'une des revendications 1 à 6, qui comprend, en outre, une huile et/ou un agent collant.
8. Formulation de bande pour administration percutanée contenant du fentanyl selon l'une des revendications 1 à 7, qui comprend, en outre, un rehausseur d'absorption transdermique.
9. Formulation de bande pour administration percutanée contenant du fentanyl selon l'une des revendications 1 à 8, dans laquelle l'adhésif autocollant comprend deux composants de polyisobutylène et de copolymère bloc styrène-isoprène-styrène.